



## BIOLOGY, ECOLOGY AND IPM OF FALL ARMY WORM *SPODOPTERA FRUGIPERDA* (J E SMITH): A REVIEW

ARFA SAFDER<sup>1</sup>, ABDUL AZIZ BUKERO<sup>2</sup>, MOAZAM HYDER<sup>3</sup>, SAMREEN RANA<sup>1</sup>, ANEESA SOHAIL<sup>1</sup>,  
MAHREEN HANIF<sup>4</sup>, SYED ANIKA SHAMSHAR<sup>5</sup>, ATTIA IMAM<sup>6</sup> AND MUHAMMAD RAMZAN<sup>7\*</sup>

<sup>1</sup>Institute of Molecular Biology and Biotechnology, The University of Lahore, 54000, Pakistan

<sup>2</sup>Sindh Agriculture University, Department of Entomology, Tandojam, Pakistan

<sup>3</sup>Key laboratory of Natural Enemy Insects, Ministry of Agriculture and Rural Affairs, Institute of Plant Protection, Chinese Academy of Agricultural Sciences, Beijing, 100193, P R China

<sup>4</sup>Institute of Plant Protection, MNS-University of Agriculture, Multan, Pakistan

<sup>5</sup>Government Graduate College (B), Vehova, Dera Ghazi Khan, Punjab, Pakistan

<sup>6</sup>Department of Plant and Environmental Protection,

National Agricultural Research Center, Islamabad, Pakistan

<sup>7</sup>State Key Laboratory for Biology of Plant Diseases and Insect Pests, Institute of Plant Protection, Chinese Academy of Agricultural Sciences, Beijing, 100193, China

\*Email: ramzan.mnsua@gmail.com (corresponding author): ORCID ID 0000-0001-8008-351X

### ABSTRACT

The fall armyworm, *Spodoptera frugiperda* (J E Smith) is an emerging polyphagous pest. Its primary food is the corn cob, which suffers significant damage with its larvae skeletonizing or “windowing” the leaves. Only the larval stage is seriously risky and small and large communities as well as researchers have used a variety of control methods. A number of low-cost mitigation strategies are available and these strategies include encouraging smallholder farmers to plant crops directly after major rainstorms, using cultural methods, such as intercropping and crop rotation, and avoiding some of the ineffective insecticides. Understanding the function of natural enemies in agricultural systems requires monitoring them in the field. *Chelonus insularis*, *Pristomerus spinator*, *Cotesia marginiventris*, *Meteorus laphygmae*, *Chelonus sonorensis*, *Orius insidiosus*, *Chelonus texanus*, *Archytas marmoratus*, *Elenomus remus*, and *Trichogramma* spp. are parasitoids of *S. frugiperda*, *Metarhizium anisopliae*, *Erynia radicans*, *Fusarium solani*, *Nomuraea rileyi* and *Beauveria bassiana* are the main entomopathogenic fungi. These biocontrol agents should be promoted to manage this pest in a sustainable manner.

**Key words:** *Spodoptera frugiperda*, maize, biology, ecology, cultural control, biological control, botanical and chemical, IPM

The fall armyworm (FAW)—*Spodoptera frugiperda* (J E Smith) is a destructive polyphagous pest and its larval stage attack all the growth stages. It is mostly found in maize growing areas of the tropical, subtropical and temperate regions (Prasanna et al., 2021; Montezano et al., 2018; Overton et al., 2021). It is native to Americas but now widely distributed. The control strategies to minimize this are becoming challenging due to various reasons such as high migratory power, morbidity and fecundity (Montezano et al., 2019; Sharanabasappa et al., 2018). This review discusses the biology, ecology, and IPM strategies. This aims to provide an understanding and basic information about the pest.

### Distribution

It is native to American continents till 2015 and cause

huge damage in many temperate areas (Rwomushana, 2019). This pest was detected in West and Central Africa in 2016 by Goergen et al. (2016). Several management strategies were used against this pest in Africa in 2016, but in few years, it invaded many African areas. In India, it was reported in 2018, while it was reported from maize in Pakistan (Naeem-Ullah et al., 2019), China (He et al., 2021), and Sri Lanka in 2019. Now it has spread to Australia, Turkey and Korea and many other countries. (Lee et al., 2020; Maino et al., 2021) and thus observed in >47 countries of Africa and 18 countries of Asia including Pakistan (Naharki et al., 2020; Rwomushana et al., 2018).

### Host plants

Being a polyphagous pest, *S. frugiperda* invade more than 353 host plants especially Poaceae, Fabaceae,

and Asteraceae families. According to the report of Montezano et al. (2018), *S. frugiperda* attack more than 76 crop families. 106 and 31 host taxa belonged to Poaceae and Asteraceae and Fabaceae families, respectively. The larvae consumed both vegetative and reproductive parts of the cultivated and wild host plant. Due to vigorous feeding, high migratory potential and fecundity, pest feed on other hosts (Sokame et al., 2020; Prasanna et al., 2021). The major host plants are maize, rice, sorghum, cotton, sunflower, lettuce, aster, safflower, barley, rye, sugarcane, wheat, millet, peanut, chickpea, soybean, alfalfa, sweet clover, garden pea, kidney bean, cowpea, red and white clover etc. (Montezano et al., 2018; Bateman et al., 2018).

### Biology

Sharanabasappa et al. (2018) observed its life history in June- July 2018. Eggs are laid in groups and a female can lay 100-200 eggs on the dorsal or ventral side of the leaves (Prasanna et al., 2018; CABI, 2018b). The eggs have a dome like shape, 0.4 and 0.3 mm dia and length respectively (Prasanna et al., 2018), while colour of eggs was creamy white as reported by Bajracharya and Bhat (2019). The 1<sup>st</sup>-2<sup>nd</sup> instar larva were green, while dark brown in 3<sup>rd</sup> to 6<sup>th</sup> instars (CABI, 2018b). The main difference from other *Spodoptera* spp., is the presence of inverted Y on the front of larval head (Prasanna et al., 2018; Ramzan et al., 2021) with four square shaped dark spots on the 8<sup>th</sup> abdominal segment. The total developmental period of larvae and pupae 14-19 and 9- 12 days, respectively on maize. The pupal period was 8.54 days as reported by Débora et al. (2017).

The forewings of male are grey and brown with triangular white patch at the apical region. A circular spot is found at the center of the forewings. Females are with greyish brown forewings; hind wings of both sexes have silver-white with a narrow dark border. The preovipositional, ovipositional and post ovipositional period was 3 to 4, 2 to 3 and 4 to 5 days, respectively. It has been reported that females live longer (Sharma et al., 2022; Srikanth et al., 2018; Tendeng et al., 2019).

### IPM strategies

The IPM strategies used against insect pests include cultural, mechanical, physical biological, botanical, microbial and chemical control. All these strategies have been used against several insect pests.

Globally, there are many cultural and physical recommended, such as timely sowing, intercropping, crop rotation (Assefa, 2018; Kasoma et al., 2020),

proper application of nitrogenous fertilizers etc. The pest population can be reduced by planting the early maturing varieties (Chhetri and Acharya, 2019). Rainfall at early stage of crop can reduce the population as reported by Harrison et al. (2019), and 56 and 70% reduction in Ghana and Zambia, respectively had been reported (Rwomushana et al., 2018). Another method is the use of traditional method i.e. crop rotation, but this cannot directly reduce the pest population due to polyphagous nature (Montezano et al., 2018) and high migratory potential (Kansiime et al., 2019). Crop rotation not only controls the pest but also prove good in improving the soil fertility that supports healthy plant growth (Prasanna et al., 2018; Meagher et al., 2016). Many plants used in push-pull technique, may lead to anti-appetizing action towards several pests like FAW (Hruska, 2019). Such types of plants can attract or repel the insect pests which provide protection to plants from pest damage (Khan et al., 2016; Hassanali, 2008). The immature stages such as larvae and egg masses can be hand-picked and this is not only adopted in Africa but as old method tested by several farmers and researchers (Yigezu and Wakgari, 2020). Many non-chemical elements such as salts, oil, urine, soaps and detergents are applied on FAW to kill them. To control the high infestation and to reduce crop losses, farmers used sand with chemicals in the form of granules (FAO, 2017). The light traps can be used to attract adults (Shimoda and Honda, 2013).

**Botanicals:** The chemicals extracted from plants have some toxic effects on insect pests. These kill the pests, reduce the larval performance and reduces the growth and development. By the application of such chemicals, the fertility and fecundity can be reduced (Rioba and Stevenson, 2020). There are several plant extracts tested against FAW but among them *Azadirachta indica*, *Phytolacca dodecandra* and *Schinus molle* are the most widely (Lin et al., 2020).

**Biological control:** Predators, parasitoids and pathogens are the main component of biological control (Nafiu et al., 2014). More than 150 parasitoids and nematodes attack the different stages of FAW (Sisay et al., 2018; Tendeng et al., 2019), while several insect and spider predators (Koffi et al., 2020) and pathogenic bacteria, fungi and viruses also invade the pest stages (Shylesha et al., 2018) which causes up to 42% mortality (Day et al., 2017; Caniço et al., 2020; Mohamed et al., 2021). *Cotesia icipe*, *Apanteles marginiventris*, *Anatrichus erinaceus*, *Chelonus insularis*, *Telenomus remus*, *Rogus laphygmae*, *Trichogramma pretiosum*, *Camponotus grioti*, *Ephisoma vitticole*, and *Meteorus*

*autographae* are the most important parasitoids as reported by Mohamed et al. (2021); Abbas et al. (2022) and Carneiro et al. (2012). Some predators such as *Pheidole megacephaly*, *Haematochares obscuripennis* and *Peprius nodulipes* had been reported by Kenis et al. (2019).

More than 750-800 fungal species are found of which *Beauveria bassiana* is the effective one (Khan and Ahmad, 2015). *Nomuraea rileyi* is another fungal species that gives best control (Romero-Arenas et al., 2014). More than 72.5% larval mortality was recorded by the use of such fungi. Another pathogen is gram positive bacteria, *Bacillus thuringiensis* which was used widely. The biological and morphological parameters of the pest have been reduced by the application of such pathogens (Farias et al., 2014; Dangal and Huang, 2015). Viruses and nematodes have also been used (Gichuhi et al., 2020; Botha et al., 2019). Entomopathogenic nematodes (EPNs) such as Steinernematidae and Heterorhabditidae are linked with some symbiotic bacteria (Sree and Varma, 2015). These have proved most effective with 100% larval mortality in combination with insecticides (Negrisoli et al., 2013); *H. indica* and *S. carpocapsae* and *S. riobravus* caused 28% and 53% larval mortality, respectively (Negrisoli et al., 2010).

**Chemical control:** The insecticides are the last option to control *S. frugiperda* as these gives quick response and protect the crop (Togola et al., 2018). Different insecticides are used of which most important are deltamethrin, cypermethrin, lambda-cyhalothrin, chlorpyrifos and permethrin. The excessive applications of these has caused environmental pollution and insecticides resistance (Sellami et al., 2015). Azevedo et al. (2004) reported that pest population can be minimized by treating seeds with some chemicals (malathion, lambda-cyhalothrin, chlorpyrifos, permethrin), Thiodicarb and clothianidin tested by Camillo et al. (2005) proved efficient on larvae. Niaz et al. (2018) concluded that indoxacarb, emamectin benzoate, methomyl, lufenuron and fubendiamide caused with maximum larval mortality.

## CONCLUSIONS

Recently, FAW has evolved into an invasive species and adapted to the environmental settings of numerous locations. Since its wide area expansion across the world, there is an urgent need to devise effective IPM strategies. It is also necessary to raise awareness among farming communities about pest

scouting, recognizing when pests attack which stage of the crop, applying treatment at the proper time, and implementing low-cost agronomic measures. Additional research should be done to examine the host plants, life cycle, and effectiveness of various control strategies. The best technique for enhancing natural enemies is to increase the population or diversity of plants. The optimum management choice for FAW may be a push-pull strategy, such as the cultivation of Napier and Desmodium grass with maize crop. The greatest method for eradicating this pest would be area-wise management and the promotion of biological fauna.

## REFERENCES

- Abbas A, Ullah F, Hafeez M, Han X, Dara M Z N, Gul H, Zhao C R. 2022. Biological Control of Fall Armyworm, *Spodoptera frugiperda*. *Agronomy* 12: 2704.
- Assefa F. 2018. Status of fall armyworm (*Spodoptera frugiperda*), biology and control measures on maize crop in Ethiopia: a review. *International Journal of Entomology and Research* 06(02): 75- 85.
- Azevedo R D, Grutzmacher A D, Loeck A E, da Silva F F, Storch G, Herpich M I. 2004. Effect of seed treatment and leaf spray of insecticides in different water volumes, on the control of *Spodoptera frugiperda* (JE Smith, 1797) (Lepidoptera: Noctuidae), in lowland corn and sorghum crops. *Revista Brasileira de Agrociencia (Brazil)* 10(1): 71-77.
- Bateman M L, Day R K, Luke B. 2018. Assessment of potential biopesticide options for managing fall armyworm (*Spodoptera frugiperda*) in Africa. *Journal of Applied Entomology* 142: 805-819.
- Botha A S, Erasmus A, du Plessis H, Van den Berg J. 2019. Efficacy of Bt Maize for Control of *Spodoptera frugiperda* (Lepidoptera: Noctuidae) in South Africa. *Journal of Economic Entomology* 112(3): 1260-1266.
- Camillo M F, Di Oliveira J R G, de Bueno A F, Bueno R C O D F. 2005. Seeds treatment on maize for *Spodoptera frugiperda* control. *Ecosistema* 30(1/2): 59-63.
- Canico A, Mexia A, Santos L. 2020. First report of native parasitoids of fall armyworm *Spodoptera frugiperda* Smith (Lepidoptera: Noctuidae) in Mozambique. *Insects* 11: 615.
- Carneiro T R, Fernandes O A. 2012. Interspecific interaction between *Telenomusremus* (Hymenoptera: Platygasteridae) and *Trichogramma pretiosum* (Hymenoptera: Trichogrammatidae) on *Spodoptera frugiperda* (Lepidoptera: Noctuidae) eggs. *Annals Academy of Bras. Ciênc.* 84: 1127-1135
- Chhetri L B, Acharya B. 2019. Fall armyworm (*Spodoptera frugiperda*): a threat to food security for south Asian country: control and management options: a review. *Farming Management* 4(1): 38-44,
- Dangal V, Huang F. 2015. Costs of Cry1F resistance in two populations of fall armyworm, *Spodoptera frugiperda* (J.E. Smith), collected from Puerto Rico and Florida. *Journal of Invertebrate Pathology* 127: 81-86.
- Day R, Abrahams P, Bateman M, Beale T, Clotley V, Cock M, Colmenarez Y, Corniani N, Early R, Godwin J. 2017. Fall armyworm: Impacts and implications for Africa. *Outlooks Pest Management* 28: 196-201.
- Débora M D S, Adeney D F B, Karine A, Cristiane D S S, Pedro M O J N, Maria C N D O. 2017. Biology and nutrition of *Spodoptera frugiperda* (Lepidoptera: Noctuidae) fed on different food sources, *Scientia Agricola* 74 (1): 18- 31.

- Farias J R, Andow D A, Horikoshi R J, Sorgatto R J, dos Santos A C, Omoto C. 2014. Field-evolved resistance to Cry1F maize by *Spodoptera frugiperda* (Lepidoptera: Noctuidae) in Brazil. *Crop Protection* 64: 150-158.
- Gichuhi J, Sevgan S, Khamis F, Van den Berg J, du Plessis H, Ekesi S, Herren J K. 2020. Diversity of fall armyworm, *Spodoptera frugiperda* and their gut bacterial community in Kenya. *Peer J* 8:e8701 <https://doi.org/10.7717/peerj.8701>
- Goergen G, Kumar P L, Sankung S B, Togola A, Tamò M. 2016. First report of outbreaks of the fall armyworm *Spodoptera frugiperda* (J E Smith) (Lepidoptera, Noctuidae), a new alien invasive pest in West and Central Africa. *PLoS ONE* 11(10): e0165632.
- Harrison R D, Thierfelder C, Baudron F, Chinwada P, Midega C, Schaffner U, Van den Berg J. 2019. Agro-ecological options for fall armyworm *Spodoptera frugiperda* (J E Smith) management: providing low-cost, smallholder friendly solutions to an invasive pest. *Journal of Environment Management*. 243: 318-330.
- Hassanali A. 2008. Integrated pest management: the push-pull approach for controlling insect pests and weeds of cereals, and its potential for other agricultural systems including animal husbandry. *Philosophical Transactions of The Royal Society of London Series B-Biologic* 363: 611-621.
- He L, Zhao S, Ali A, Ge S, Wu K. 2021. Ambient humidity affects development, survival, and reproduction of the invasive fall armyworm, *Spodoptera frugiperda* (Lepidoptera: Noctuidae) in China. *Journal of Economic Entomology* 114(3): 1145- 1158.
- Hruska A J. 2019. Fall armyworm (*Spodoptera frugiperda*) management by smallholders. *CAB Review* 14(043): 1-11.
- Kansiime M K, Mugambi I, Rwomushana I, Nunda W, Lamontagne-Godwin J, Rware H, Phiri N A, Chipabika G, Ndlovu M, Day R. 2019. Farmer perception of fall armyworm (*Spodoptera frugiperda* J.E. Smith) and farmlevel management practices in Zambia. *Pest Management Science* 75, 2840-2850.
- Kasoma C, Shimelis H, Laing M D. 2020. Fall armyworm invasion in Africa: implications for maize production and breeding. *Journal of Crop Improvement* 35(1): 111-146.
- Kenis M, DuPlessis H, VandenBerg J, Ba M N, Goergen G, Kwadjo K E, Baoua I, Tefera T, Buddie A, Cafà G. 2019. *Telenomusremus*, a Candidate Parasitoid for the Biological Control of *Spodoptera frugiperda* in Africa, is already Present on the continent. *Insects* 10: 92-103.
- Khan Z R, Midega C A O, Hooper A M, Pickett J A. 2016. Push-pull: chemical ecology-based integrated pest management technology. *Journal of Chemistry and Ecology* 42: 689-697.
- Koffi D, Kyerematen R, Eziah V Y, Agboka K, Adom M, Goergen G, Meagher R L. 2020. Natural enemies of the fall armyworm, *Spodoptera frugiperda* (J.E. Smith) (Lepidoptera: Noctuidae) in Ghana. *Florida Entomologist* 103: 85
- Lee G S, Seo B Y, Lee J, Kim H, Song J H, Lee W. 2020. First Report of the fall armyworm, *Spodoptera frugiperda* (Smith, 1797) (Lepidoptera, Noctuidae), a new migratory pest in Korea. *Korean Journal of Applied Entomology* 59(1): 73-78
- Lin S K, Liu K H, Wang R F, Liu B J, Zhang Y, Wu J Y Z, Cheng D M, Xu H H, Zhang Z X. 2020. Indoor toxicity of azadirachtin to *Spodoptera frugiperda* and its control effect in field. *Journal of South China Agricultural University* 41: 22-27.
- Maino J L, Schouten R, Overton K, Day R, Ekesi S, Bett B, Barton M, Gregg P C, Umina P A, Reynolds O L. 2021. Regional and seasonal activity predictions for fall armyworm in Australia. *Current Research in Insect Science* 1: 100010.
- Meagher R L, Nuessly G S, Nagoshi R N, Hay-Roe M M. 2016. Parasitoids attacking fall armyworm (Lepidoptera: Noctuidae) in sweet corn habitats. *Biology Control* 95: 66- 72.
- Mohamed S A, Wamalwa M, Obala F, Tonnang HEZ, Tefera T, Calatayud P A. 2021. A deadly encounter: Alien invasive *Spodoptera frugiperda* in Africa and indigenous natural enemy, *Cotesia icipe* (Hymenoptera, Braconidae). *PLoS ONE* 16(7): e0253122
- Montezano D G, Specht A, Sosa-Gómez D R, Roque-Specht V F, Malaquias J V, Paula-Moraes S V, Hunt T E. 2019. Biotic potential and reproductive parameters of *Spodoptera frugiperda* (J.E. Smith, 1797) (Lepidoptera: Noctuidae). *Journal of Agricultural Science* 11(13): 240-252.
- Montezano D G, Specht A, Sosa-Gómez D R, Roque-Specht V F, Sousa-Silva J C, Paula-Moraes S D, Hunt T E. 2018. Host plants of *Spodoptera frugiperda* (Lepidoptera: Noctuidae) in the Americas. *African Entomology* 26(2):286-300. <https://doi.org/10.4001/003.026.0286>
- Nafiu B S, Dong H, Mustapha S. 2014. Biological control of insect pests in West Africa: a review. *International Journal of Applied Research Technology* 3(9): 39-45, [doi.org/10.13140/RG.2.2.13034.31687](https://doi.org/10.13140/RG.2.2.13034.31687)
- Negrisoni A S, Garcia M S, Negrisoni C R C B. 2010. Compatibility of entomopathogenic nematodes (Nematoda: Rhabditida) with registered insecticides for *Spodoptera frugiperda* (Smith, 1797) (Lepidoptera: Noctuidae) under laboratory conditions. *Crop Protection* 29: 545-549.
- Negrisoni C R C B, Negrisoni A S, Botton M, Garcia M S, Bernardi D. 2013. Evaluation of efficacy of 18 strains of entomopathogenic nematodes (Rhabditida) against *Planococcus citri* (Risso, 1813) (Hemiptera: Pseudococcidae) under laboratory conditions. *Experimental Parasitology* 134(3): 295-298.
- Niaz U, Khan A, Javed M U. 2018. Fall armyworm *Spodoptera frugiperda* (Lepidoptera: Noctuidae) a well-known quarantine pest and its global IPM strategies. *Journal of Agricultural Science and Botany* 2(3): 1-4.
- Overton K, Maino J L, Day R, Umina P A, Bett B, Carnovale D, Reynolds O L. 2021. Global crop impacts, yield losses and action thresholds for fall armyworm (*Spodoptera frugiperda*): A review. *Crop Protection* (Guildford, Surrey) 145: 105641.
- Prasanna B M, Huesing J E, Peschke V M, Eddy R. 2021. Fall Armyworm in Asia: A Guide for Integrated Pest Management. Mexico, CDMX: CIMMYT.
- Ramzan M, Ilahi H, Adnan M, Ullah A, Ullah A. 2021. Observation on fall armyworm, *Spodoptera frugiperda* (Lepidoptera: Noctuidae) on maize under laboratory conditions. *Egyptian Academic Journal of Biological Sciences* 14: 99-104.
- Ramzan M, Usman M, Sajid Z, Ghani U, Basit M A, Razaq M, Shafee W, Shahid M R. 2020. Bio-ecology and management of *Spodoptera frugiperda* (Lepidoptera: Noctuidae): A review. *Journal of Pure and Applied Agriculture* 5(4):1-9.
- Rioba N B, Stevenson P C. 2020. Opportunities and scope for botanical extracts and products for the management of fall armyworm (*Spodoptera frugiperda*) for smallholders in Africa. *Plants*, 9, 207.
- Romero-Arenas O, Rivera A, Aragon A, Parraguire C, Cabrera E, Lopez F. 2014. Mortality Evaluation of Armyworm (*Spodoptera frugiperda* J. E. Smith) by using *Metarhizium anisopliae* In vitro. *Journal of Pure and Applied Microbiology* 8(2):59-67.
- Rwomushana I, Bateman M, Beale T, Beshe P, Cameron K, Chiluba M, Tambo J. 2018. Fall armyworm: impacts and implications for Africa: evidence note update UK: CABI.
- Rwomushana I. 2019. *Spodoptera frugiperda* (fall armyworm). *Invasive Species Compendium*. Wallingford, UK: CABI. <https://doi.org/10.1079/ISC.29810.20203373913>

- Sellami S, Mesrati A, Tounsi S, Jamoussi K. 2015. Toxicity, activation process, and histopathological effect of *Bacillus thuringiensis* vegetative insecticidal protein Vip3Aa16 on *Tuta absoluta*. *Applied Biochemistry and Biotechnology* 175:1992-1999
- Sharanabasappa, Kalleshwaraswamy C M, Asokan R, Mahadevaswamy H M, Maruthi M S, Pavithro H B, Kavita H, Shivaray N, Prabhu S T, Georg G. 2018. First report of fall army worm, *Spodoptera frugiperda* (J.E. Smith) (Lepidoptera: Noctuidae) an alien invasive pest on maize in India. *Pest Management in Horticultural Ecosystems* 24(1): 23-29.
- Sharma S, Tiwari S, Thapa R B, Neupane S, Reddy G V P, Pokhrel S, Muniappan R. 2022. Life cycle and morphometrics of fall armyworm (*Spodoptera frugiperda*) (Lepidoptera: Noctuidae) on maize crop. *SAARC Journal of Agriculture*. 20(1): 77-86.
- Shimoda M, Honda K. 2013. Insect reactions to light and its applications to pest management. *Applied Entomology Zoology* 48: 413-421.
- Shylesha A N, Jalali S K, Gupta A, Varshney R, Venkatesan T, Shetty P, Ojha R, Ganiger P C, Navik O, Subaharan K, Bakthavatsalam N. 2018. Studies on new invasive pest *Spodoptera frugiperda* (JE Smith)(Lepidoptera: Noctuidae) and its natural enemies. *Journal of Biological Control* 32(3): 145-51.
- Sisay B, Tefera T, Wakgari M, Ayalew G, Mendesil E. 2019. The efficacy of selected synthetic insecticides and botanicals against fall armyworm, *Spodoptera frugiperda*, in Maize. *Insects* 10: 45.
- Sree K S, Varma A. 2015. Biocontrol of lepidopteran pests. Switzerland: Springer International Publishing <https://doi.org/10.1007/978-3-319-14499-3>
- Srikanth J, Geetha N, Singaravelu B, Ramasubramanian T, Mahesh P, Saravanan L, Salin K P, Chitra N, Muthukumar M. 2018. First report of occurrence of fall armyworm *Spodoptera frugiperda* in sugarcane from Tamil Nadu, India. *Journal of Sugarcane Research* 8: 195-202.
- Tendeng E, Labou B, Diatte M, Djiba S, Diarra K. 2019. The fall armyworm *Spodoptera frugiperda* (JE Smith), a new pest of maize in Africa: biology and first native natural enemies detected. *International Journal of Biological and Chemical Sciences* 13: 1011-1026.
- Tendeng E, Labou B, Diatte M, Djiba S, Diarra K. 2019. The fall armyworm *Spodoptera frugiperda* (J.E. Smith), a new pest of maize in Africa: Biology and first native natural enemies detected. *International Journal of Biology Chemistry Science* 13: 1011-1026.
- Togola A, Meseka S, Menkir A, Badu-Apraku B, Boukar O, Tamo M, Djouaka R. 2018. Measurement of pesticide residues from chemical control of the invasive *Spodoptera frugiperda* (Lepidoptera: Noctuidae) in a maize experimental field in Mokwa, Nigeria. *International Journal of Environmental Research and Public Health* 15(5): 849.
- Yigezu G, Wakgari M. 2020. Local and indigenous knowledge of farmers management practice against fall armyworm (*Spodoptera frugiperda*) (J. E. Smith) (Lepidoptera: Noctuidae): a review. *Journal of Entomology and Zoology Studies* 8(1): 765-770.

(Manuscript Received: January, 2023; Revised: February, 2023;

Accepted: February, 2023; Online Published: February, 2023)

Online First in [www.entosocindia.org](http://www.entosocindia.org) and [indiantentomology.org](http://indiantentomology.org) Ref. No. e23027