

Indian Journal of Entomology Online published Ref. No. e23234

FORTUITOUS INTRODUCTION OF LANTANOPHAGA PUSILLIDACTYLUS (WALKER) INTO NEPAL

R. MUNIAPPAN¹* AND LALIT SAH²

¹CIRED, Virginia Tech, 526 Prices Fork Road, Blacksburg, VA 24061, USA ²Feed the Future Nepal Integrated Pest Management, iDE, Lalitpur, Nepal *Email: rmuni@vt.edu (corresponding author): ORCID ID: 0000-0002-2992-2792

ABSTRACT

The neotropical plant, *Lantana camara* L., introduced as an ornamental plant to several countries, has become an invasive weed. It was first reported from Nepal in 1848. About 50 countries have introduced natural enemies for suppression of the weed. India, Nepal's neighboring country, initiated biological control of this weed in 1916 and reported the occurrence of *Lantanophaga pusillidactylus* Walker. In a recent survey, the Feed the future Nepal Integrated Pest Management project found larvae of this insect feeding on flower heads of *L. camara* in Nepal. This communication reports the first discovery of *L. pusillidactylus* in Nepal on March 13, 2023.

Key words: Lantana camara, Lantanophaga pusillidactylus, fortuitous introduction, biological control of a weed, Invasive species management

The alien invasive weed, Lantana camara L. (Verbanaceae) is neotropical in origin and pantropical in distribution. It invades pastures, orchards, plantations, forests, vacant lands, and roadsides and it has established in about 70 countries worldwide (Day et al., 2003). It was introduced into the Royal Botanic Garden in Calcutta (now known as the Acharya Jagadish Chandra Bose Botanic Garden, Kolkata), India in 1809 as an ornamental plant, and subsequently introduced in other places. Since then, it has escaped cultivation and become a serious weed (Rao et al., 1971). It was first reported from Nepal in 1848, and, currently, is distributed in the elevations from 70 to 1715 (masl) (Shrestha, 2016). Lantana camara is a composite species. While it is thought to have evolved from two or more species, over 650 varieties are known to exist (Day et al., 2003). It grows as individual clumps or dense thickets, displacing desirable species (Day and Zalucki, 2009). It is highly allelopathic, reduces productivity of orchards, and pastures and interferes with the movement of wildlife in forests. Options available to manage this weed include use of chemical herbicides, mechanical removal, use of fire, adoption of cultural methods, and release of biological agents. Of these, biological control is the most preferred long-time option considering its establishment in forests, orchards, pastures, fences, vacant lots, slopes and remote areas.

Hawaii was first to import and introduce natural enemies of *L. camara* from Mexico in 1902. Since then, 41 agents have been released in 50 countries resulting

only in partial suppression and inadequate control (Julien and Griffiths, 1998; Day et al., 2003). From November 15, 1916 to March 31, 1919, Ramachandra Rao conducted a survey of arthropods recruited by L. camara in India and Burma (now Myanmar). He recorded 148 insects, two pathogenic fungi and one parasitic plant associated with L. camara (Ramachandra Rao, 1920). Of these, occurrence of one, a flower feeder Lantanophaga (Platyptilia) pusillidactylus (Walker) (Lepidoptera: Pterophoridae), and a native of Mexico and the Caribbean region, was recorded in the survey. Most likely it was fortuitously introduced to India with the imported L. camara plant material. In fact, L. pusillidactylus was one of the insects imported by the Hawaii Department of Agriculture from Mexico for control of L. camara in 1902 (Perkins and Swezey, 1924). From Hawaii, it was introduced to Pohnpei in 1948 and from there to Palau in 1960. Since then, it has fortuitously established in several Micronesian Islands (Denton et al., 1991; Muniappan, 1988; 1989; Muniappan and Reddy, 2003). Currently L. pusillidactylus has established in Australia, Federated States of Micronesia, Guam, Hawaii, New Zealand, Northern Mariana Islands, Palau, Papua New Guinea, and Solomon Islands in Oceania; Bhutan, China, Hongkong, India, Indonesia, Israel, Myanmar, Philippines, Sri Lanka, Taiwan, Thailand, and Vietnam in Asia; Canary Islands, Cape Verde, Democratic Republic of Congo, Republic of Congo, Egypt, Eswatini, Ivory Coast, Madagascar, Madeira, Malawi, Mauritius, Morocco, Namibia, Nigeria, Reunion, Saint Helena, Seychelles, South Africa, Tanzania, Zambia, and Zimbabwe in Africa; and Cyprus, Italy, Malta, and Portugal in Europe; and Argentina, Barbados, Bermuda, Brazil, Cuba, Dominican Republic, Ecuador, Galapagos, Grenada, Guiana, Jamaica, Mexico, Paraguay, Peru, Puerto Rico, Surinam, Virgin Islands, and U.S.A. in the Americas (Agius, 2017; Day et al., 2003; Day and Zalucki, 2009; De Prins and de Prins, 2017); Ramachandra Rao, 1920). This moth is one of the few lantana agents that can tolerate wide climatic conditions (Day et al., 2003).

On March 13, 2023, the Feed the Future Nepal Integrated Pest Management (FTFNIPM) project found *L. pusillidactylus* on *L. camara* at Mehelkuna, Gurvakot municipality, Surkhet district, Karnali province, Nepal (28.43675227422874, 81.84756356309164) (Fig. 1). It lays individual eggs on flower buds and the eggs hatch in four days. Larvae bore in to the flower buds and each larva can destroy at least 15 to 20 flowers in a flower head as a result the flower head carries only 3 or 4 berries instead of a large bunch. Pupation takes place in a silken cocoon in the flower head. In Guam, Muniappan (1988) reported it to infest 32% to 83% flower heads in different parts of the island. In his study from 1916 to 1919, Ramachandra Rao (1920) reported *L. pusillidactylus* as one of the most efficient of the insects on lantana in India. He also noted that one egg parasitoid, some larval parasitoids and a predator of the pupa *Icaria* sp. (Hymenoptera: Vespidae) affect the efficacy of this insect.

This is a first report of a fortuitous introduction of *L. pusillidactylus* on *L. camara* in Nepal. Earlier reports of such fortuitous introductions include natural enemies of the weeds *Parthenium hysterophorus* L. (Asteraceae) and *Ageratina adenophora* (Spreng.) R.M. King & H. Robinson (Asteraceae). The leaf feeding beetle *Zygogramma bicolorata* Pallister (Coleoptera: Chrysomelidae) and winter rust *Puccinia abrupta* Diet. & Holw. var. *partheniicola* (Jackson) (Pucciniales: Pucciniaceae) of *P. hysterophorus* and the stem galling fly *Procecidochares utilis* Stone (Diptera: Tephritidae) and the leaf spot fungus *Passalora ageratinae* Crous and A.R. Wood (Capnoidales: Mycosphaerellaceae) of *A. adenophora* were introduced (Shrestha, 2016).

Even though *L. camara* has been recognized as an exotic invasive weed in forests and shrublands in Nepal (Shrestha, 2016), no control method has been adopted except manual digging out the clumps in a few cases. *Lantanophaga pusillidactylus* is the first natural enemy of *L. camara* to be reported from Nepal. A systematic



Fig. 1. Location of first finding of L. pusillidactylus in Nepal

survey of natural enemies recruited by *L. camara* in Nepal would form a baseline study for any future biocontrol activities to be taken up.

ACKNOWLEDGEMENTS

The authors are grateful to team members of International Development Enterprises (iDE) and Catholic Relief Services (CRS) who accompanied us in the field trips.

FINANCIAL SUPPORT

Funding support provided by the USAID mission in Nepal Award Number 72036720LA00001 and CRS Farmer to Farmer project NE2119.

AUTHORS CONTRIBUTION STATEMENT

R M conceived and R M and L S conducted field surveys. R M wrote the manuscript. All authors read and approved the manuscript.

CONFLICT OF INTEREST

Authors declare there is no conflict or competing interests.

REFERENCES

Agius J. 2017. *Lantanophaga pusillidactylus* (Walker, 1864) new to the Maltese Islands (Lepidoptera: Pterophoridae). SHILAP Revista de Lepidopterologia 45: 259-261

Day M D, Wiley V J, Playford J, Zalucki M P. 2003. Lantana: Current

(Manuscript Received: April, 2023; Revised: April, 2023; Accepted: April, 2023; Online Published: April, 2023) Online First in www.entosocindia.org and indianentomology.org Ref. No. e23234

Management Status and Future Prospects. Canberra, Australia: ACIAR Monograph Series. 129 p.

- Day M D, Zalucki M P. 2009. Lantana camara Linn. (Verbenaceae). In, Muniappan R, Reddy G V P, Raman A. (eds.). Biological Control of Tropical Weeds using Arthropods. Cambridge University Press. pp. 211-246.
- Denton G R W, Muniappan R, Marutani M. 1991. The distribution and biological control of *Lantana camara* in Micronesia. Micronesia Supplement 3: 71-81.
- De Prins J, De Prins W. 2017. *Lantanophaga pusillidactylus* (Walker, 1864). Afromoths. Retrieved November 18, 2017.
- Julien M H, Griffiths M W. 1998. Biological Control of Weeds. A World Catalogue of Agents and their Target Weeds. Fourth Edition. CABI Publishing, Wallingford, UK.
- Muniappan R. 1988. Biological control of the weed, *Lantana camara* in Guam. Journal of Plant Protection in the Tropics 5: 99-101.
- Muniappan R. 1989. Biological control of *Lantana camara* L. in Yap. Proceedings, Hawaiian Entomological Society 29: 195-196.
- Muniappan R, Reddy G V P. 2003. Fortuitous introduction of two natural enemies of *Lantana camara* to Chuuk. Proceedings, Hawaiian Entomological Society 36: 123-124.
- Perkins R C, Swezey O H. 1924. The introduction into Hawaii of insects that attack lantana. Bulletin of the Hawaiian Sugar Planters' Association 16: 1-83.
- Ramachandra Rao Y. 1920. Lantana insects in India. Report on an enquiry into the efficiency of indigenous insect pests as a check on the spread of lantana in India. Memoirs of the Department of Agriculture, Indian Entomological Service 5: 239-314.
- Rao V P, Ghani M A, Sankaran T, Mathur K C. 1971. A Review of the Biological Control of Insects and Other Pests in South-East Asia and the Pacific Region. Commonwealth Agricultural Bureaux, Farnham Royal, Slough SL2 3BN, England. 149 p.
- Shrestha B B. 2016. Invasive alien plant species. In, Jha P K, Siwakoti M, Rajbhandari S. (eds.). Frontiers in Botany, Central Department of Botany, Tribhuvan University, Kirtipur, Kathmandu, Nepal, 269-284.