



## POPULATION DYNAMICS OF INSECT PESTS OF PEA

SHRADDHA TARE<sup>1\*</sup>, NANDA KHANDWE<sup>1</sup> AND NAVEEN<sup>2</sup>

<sup>1</sup>Department of Entomology, RVSKVV- Rafi Ahmed Kidwai College of Agriculture, Sehore 466001, Madhya Pradesh, India

<sup>2</sup>Department of Entomology, College of Agriculture,

Rajmata Vijayaraje Scindia Vishwa Vidyalyaya, Gwalior 474002, Madhya Pradesh, India

\*Email: tareshraddha09@gmail.com (corresponding author): ORCID ID 0009-0000-7688-615X

### ABSTRACT

The seasonal incidence of major insect pests on pea *Pisum sativum* at different growth stages has been assessed in this study. Major pests observed include gram pod borer *Helicoverpa armigera* (Hubner), blue butterfly *Lampedes boeticus* (L.), pea leaf miner *Chromatomyia horticola* Goureau and pea aphid *Acyrtosiphon pisum* (Harris). These results revealed that peak incidence of *H. armigera* (7.80 larvae/ plant) and *L. boeticus* (8.60 larvae/ plant) was during the 8<sup>th</sup> and 9<sup>th</sup> SMW, respectively; while for *C. horticola* (5.50 larvae/ leaf) and *A. pisum* (94.1 aphids/ plant) it was 5<sup>th</sup> SMW. *Helicoverpa armigera* incidence revealed a significant negative correlation with minimum temperature ( $r = -0.628$ ) while for *L. boeticus* it was positive one with maximum temperature ( $r = 0.604$ ). The incidence of *C. horticola* was observed to be significantly negatively correlated with minimum temperature and rainfall ( $r = -0.826$  and  $r = -0.584$ , respectively); while *A. pisum* revealed a significant negative correlation with minimum temperature ( $r = -0.806$ ).

**Key words:** Field pea, *Pisum sativum*, insect pests, seasonal incidence, *Helicoverpa armigera*, *Lampedes boeticus*, *Chromatomyia horticola*, *Acyrtosiphon pisum*, weather parameters, correlation, pod borers

Pea *Pisum sativum* L., belonging to the family Fabaceae, is an important vegetable grown in the Indian subcontinent, in the milder temperate zones. It is cultivated on 0.64 million ha in India, with production of 0.88 mt and productivity of 1375 kg/ ha (Anonymous, 2020) It serves as a cheap source of digestible protein, carbohydrates, fat, vitamins, and minerals (Tiwari et al., 2019). Losses due to insect pests are a handicap for its maximum yield. From seedling to harvest, 24 insect pests have been reported (Bijjir and Verma, 1995). Of these, the gram pod borer *Helicoverpa armigera* (Hubner), blue butterfly *Lampides boeticus* (L.), pea pod borer *Etiella zinckenella* (Treitsshke), pea aphid *Acyrtosiphon pisum* (Harris), pea leaf miner *Phytomyza horticola* (Goureau), pod fly *Melanogromyza obtuse* (Malloch), pea stem fly, *Melanogromyza phaseoli* (Tryon) and thrips *Caliothrips indicus* (Bagnall) cause serious losses (Mittal and Ujagir, 2007; Yadav et al., 2015). The correlation studies between seasonal insect pest occurrence and succession patterns helps to understand how changing plant communities and habitats influence pest populations. For an effective IPM, knowledge on the seasonal incidence and abundance of pests, and their population dynamics is required and hence the present study.

### MATERIALS AND METHODS


A field experiment was carried out during rabi, 2020-21 at the Agricultural Research Farm, Rafi Ahmed Kidwai College of Agriculture, Sehore (23°12'N, 77°05'E). The field was prepared using standard packages and practices and a non-replicated trial was laid out in the field with a plot size of 50 m<sup>2</sup> having row to row distance of 30 cm and plant to plant distance of 10 cm, with vegetable pea variety 'Arkel'. The experimental plot was kept free from insecticidal spray, and data on seasonal incidence were recorded at weekly intervals starting from the appearance of respective pest till harvesting. For the pod borer complex, number of larvae were counted from ten randomly selected plants/ plot; number of larvae of leaf miners were recorded from 3 leaves, one each from the upper, middle, and lower canopy. The incidence of the *A. pisum* were recorded by counting their number from the tender parts of ten randomly selected plants. Weekly data on maximum and minimum temperature, relative humidity, and rainfall were obtained from the meteorological observatory of the Rafi Ahmed Kidwai College of Agriculture, Sehore. These were used to work out a simple correlation with data on incidence using MS Excel.

**RESULTS AND DISCUSSION**

During rabi, 2020-21, four major insect pests viz., *H. armigera*, *L. boeticus*, *C. horticola* and *A. pisum* were observed at different stages of growth. Larvae of *H. armigera* started to appear from the 48<sup>th</sup> Standard Meteorological Week (SMW) (29<sup>th</sup> November to 05<sup>th</sup> December) with 0.4 larvae/ plant (Fig. 1); it continued to increase from 49<sup>th</sup> to 7<sup>th</sup> SMW (2<sup>nd</sup> week of December to the 3<sup>rd</sup> week of February), which varied from 1.00 to 7.20 larvae/ plant, with peak incidence during 8<sup>th</sup> SMW (24<sup>th</sup> February) with 7.80 larvae/ plant; this continued till maturity of the crop during 11<sup>th</sup> SMW (14<sup>th</sup> to 20<sup>th</sup> March). The correlation analysis showed that there was significant negative correlation of minimum temperature with larval incidence( $r = -0.628$ ), while others were non-significant. These findings are in agreement with those of Dubey et al. (1993) on *H. armigera*, with its peak activity beginning in February and lasting until March. Prasad et al. (1997) also found that the adults of *H. armigera* reached maximum during late March. Pandey et al. (2002) observed a significant

positive linear relationship with maximum temperature (0.64), minimum temperature (0.62) and evaporation (0.60). Yadav et al. (2019) also found that *H. armigera* larval incidence starts increasing during third week of December to first week of March. *Lampides boeticus* started appearing during flowering and pod formation stage and continued up to the full maturity of the crop (Fig. 1); it first appeared during 52<sup>nd</sup> SMW (27<sup>th</sup> Decemberto 02<sup>nd</sup> January- 0.9 larvae/ plant), continued to increase from 1<sup>st</sup> to 8<sup>th</sup> SMW (1<sup>st</sup> week of January to the 4<sup>th</sup> week of February), and peak was during 9<sup>th</sup> SMW (26<sup>th</sup> February to 4<sup>th</sup> March- 8.60 larvae/ plant). Its incidence continued till maturity with 7.70 larvae/ plant during 11<sup>th</sup> SMW (14<sup>th</sup> to 20<sup>th</sup> March). A statistically significant and positive association was found between the larval incidence and the maximum temperature ( $r = 0.604$ ). These findings are partially in accordance with Kaushik and Singh (1982).

The incidence of *C. horticola* commenced during 51<sup>st</sup> SMW (4<sup>th</sup> week of December- 3.4 larvae/ leaf), and then continued to increase from 52<sup>nd</sup> to 4<sup>th</sup> SMW (27<sup>th</sup>



Insect pests	Weather parameters						
	Maximum temperature (°C)		Minimum temperature (°C)		Rainfall (mm)		Relative humidity (%)
	r	b <sub>yx</sub>	r	b <sub>yx</sub>	r	b <sub>yx</sub>	
<i>H. armigera</i>	0.357	-	-0.628*	-0.70	-0.465	-	-0.260
<i>L. boeticus</i>	0.604*	0.50	-0.382	-	-0.441	-	-0.345
<i>C. horticola</i>	-0.097	-	-0.826*	-0.73	-0.584*	-3.76	-0.055
<i>A. pisum</i>	0.061	-	-0.806*	-9.85	-0.447	-	-0.092

\*Significant at p= 0.05

Fig. 1. Population dynamics of *H. armigera*, *L. boeticus*, *C. horticola* and *A. pisum* in pea (rabi, 2020-21)

December to 30<sup>th</sup> January- 3.6 to 5.1 larvae/ leaf) with peak being during 5<sup>th</sup> SMW (4<sup>th</sup> February- 5.50 larvae/ leaf). It continued till the maturity (3.1 larvae/ leaf- 11<sup>th</sup> SMW -14<sup>th</sup> to 20<sup>th</sup> March). The incidence revealed a significant negative correlation with both minimum temperature and rainfall ( $r = -0.826$  and  $r = -0.584$ , respectively). These results are partially in accordance with Singh et al. (2013). The incidence of *A. pisum* started during 4<sup>th</sup> week of November (47<sup>th</sup> SMW- 6.5 aphids/ plant), and continued to increase from 48<sup>th</sup> to 4<sup>th</sup> SMW (29<sup>th</sup> November to 30<sup>th</sup> January- 13.9 to 87.0 aphids/ plant); and during 5<sup>th</sup> SMW (4<sup>th</sup> February) it was at its peak of 94.1 aphids/ plant, and it continued till maturity. Its incidence showed a significant negative correlation with minimum temperature ( $r = -0.806$ ). Melesse and Singh (2012) found similar results with temperature and rainfall. Patel et al. (2023) found that aphid incidence had a negative correlation with minimum and maximum temperature and morning relative humidity, while it was positive with rainfall and relative humidity.

#### ACKNOWLEDGEMENTS

The authors acknowledge the Head of the Department of Entomology, Genetics and Plant Breeding and the Dean, Rafi Ahmed Kidwai College of Agriculture, Sehore for providing facilities and for support.

#### FINANCIAL SUPPORT

No funding received.

#### AUTHOR CONTRIBUTION STATEMENT

S T and N K conceived, designed research and conducted experiments. S T and N contributed to analytical tools. S T wrote the manuscript. All authors read and approved the manuscript.

#### CONFLICT OF INTEREST

No conflict of interest.

(Manuscript Received: May, 2023; Revised: August, 2023;

Accepted: September, 2023; Online Published: October, 2023)

Online First in [www.entosocindia.org](http://www.entosocindia.org) and [indianentomology.org](http://indianentomology.org) Ref. No. e23287

#### REFERENCES

- Anonymous. 2020. Selected State-wise Area, Production and Productivity of Rabi Peas in India (2020-2021). <https://www.indiastatagri.com/table/agriculture/state-season-wise-area-production-productivity-tot/1409310>. accessed on 25/07/2021.
- Bijjur S, Verma, S. 1995. Effect of abiotic factors on the pests of pea and natural enemies. *Indian Journal of Entomology* 57(3): 233-239.
- Dubey O P, Odak S C, Gargav V P. 1993. Population dynamics of gram pod borer. *Jawaharlal Nehru Krishi Vishwa Vidyalaya Research Journal* 27(1): 59-63.
- Kaushik S K, Singh G. 1982. A short note on the incidence of blue butterfly, *Lampides boeticus* L. (Lycaenidae: Lepidoptera). *Indian Journal of Plant Protection* 10: 94.
- Mittal V, Ujagir R. 2007. Succession of insect pests associated with pea crop (*Pisum sativum* Linnaeus) at Pantnagar, India. *Environment and Ecology* 25(4): 1030-1035.
- Melesse T, Singh S K. 2012. Effect of climatic factors on pea aphid, *Acyrtosiphon pisum* Harris (Homoptera: Aphididae) population and its Management through planting dates and biopesticides in field pea (*Pisum sativum* L.). *Journal of Agricultural Technology* 8(1): 125-132.
- Prasad C S. 1997. Host range and seasonal incidence of *Helicoverpa armigera* Hubner in the lower Kumaon Hills of Uttar Pradesh, Indian. *Insect Environment* 3(2): 33-34.
- Pandey R K, Singh G R. 2002. Effect of integrated pest management technology on yield of field pea in eastern parts of Uttar Pradesh. *Proceedings. National symposium on integrated management in horticultural crops for protection of environment and biodiversity*. pp. 147-148.
- Patel D, Kumar R, Patel A, Prajapati G, Kumar, S. 2023. Seasonal incidence of insect-pests of field pea and correlation in relation to weather parameters. *Journal of Entomological Research* 47(2): 405-408.
- Singh M, Mishra T. 2013. Seasonal incidence of pea leaf miner, *Chromatomyia horticola* infesting pea. *Plant Archives* 13(2): 941-943
- Tiwari A, Markam N, Dubey M K. 2019. Factors responsible for adoption of improved pea production technology among the pea growers. *International Journal of Current Microbiology and Applied Sciences* 8(3): 933-938.
- Yadav S K, Patel S. 2015. Insect pest complex on *Pisum sativum* L. and their natural enemies at Pantnagar. *Journal of Plant Development Sciences* 7(11): 839-841.
- Yadav A, Singh V, Yadav A, Singh H. 2019. Studies on succession of insect pest complex associated with pea at Bikaner. *Journal of Entomology and Zoology Studies* 7(3):1606-1608.